



This Journal of Environmental Horticulture article is reproduced with the consent of the Horticultural Research Institute (HRI – www.hriresearch.org), which was established in 1962 as the research and development affiliate of the American Nursery & Landscape Association (ANLA – <http://www.anla.org>).

HRI's Mission:

To direct, fund, promote and communicate horticultural research, which increases the quality and value of ornamental plants, improves the productivity and profitability of the nursery and landscape industry, and protects and enhances the environment.

The use of any trade name in this article does not imply an endorsement of the equipment, product or process named, nor any criticism of any similar products that are not mentioned.

Growth Response of *Photinia* and *Thuja* and Nutrient Concentration in Tissues and Potting Medium as Influenced by Composted Sewage Sludge, Peat, Bark and Sawdust in Potting Media¹

R.L. Ticknor, D.D. Hemphill, Jr. and D.J. Flower²
North Willamette Agricultural Experiment Station
Oregon State University
Aurora, OR 97002

Abstract

Composted sewage sludge was blended with pumice and either bark, peat moss, or sawdust to form 6 potting media. Increasing the proportion of compost in media from 25 to 50% caused increased initial pH, electrical conductivity (EC), and air-filled pore space (AS) in all media, and increased growth of *Photinia x fraseri* Dress. and *Thuja occidentalis* L. 'Pyramidalis.' Sawdust-containing media had the highest pH, EC, and AS; sphagnum peat-containing media the highest water holding capacity and greatest shrinkage; bark-based media the highest bulk density. Plant growth in compost-based media containing peat or bark was equal to or better than that in two commercial media composed primarily of bark or peat.

Index words: *Photinia x fraseri*, *Thuja occidentalis* 'Pyramidalis,' peat, bark, sawdust, pumice, air-filled pore space, electrical conductivity, pH

Introduction

A wide range of components are used to produce soil-less potting media for container production of woody plants. Peat-based media have been particularly successful, but their high cost has encouraged a search for substitutes.

The physical and chemical properties of media which stimulate plant growth have been widely investigated. Critical ranges for bulk density (7,13), air-filled pore space (3,12,13), organic matter (7), cation exchange capacity (7), water holding capacity (3), and particle size distribution (9,12,13) have been determined.

Composts of sewage sludge with municipal solid waste (11) or wood chips (6,14) have been used in nursery crop production. Similar composts are becoming available as more municipalities turn to composting for waste disposal. Portland, Oregon produces a sewage sludge which contains levels of plant nutrients and trace elements near or slightly lower than the national median (4). The Portland sludge is now being composted with conifer sawdust and marketed as a nursery crop potting medium.

The purpose of this study was to determine the physical and chemical characteristics of 6 compost-amended media and to measure their effects on the growth and elemental content of 2 woody nursery crops.

Materials and Methods

Polymer dewatered, anaerobically digested municipal sewage sludge from Portland, Oregon was composted with conifer sawdust in a closed vessel pilot plant (10) by

the Taulman-Weiss Co. of Atlanta, GA. The compost was not screened to remove wood particles. Six potting mixes were prepared in a factorial combination of either 25 or 50% by volume compost, 50 or 25% sphagnum peat moss, conifer sawdust, or hammer-milled conifer bark, and 25% pumice amended with 1.77 kg gypsum/m³ (3 lb/yd³). Physical and chemical characteristics of the resulting media are shown in Table 1. Plants growing in these media were compared to plants in 2 commercial media. Commercial medium A consisted of 1.3 cm (½ in) minus conifer bark supplemented with 6.5 kg of 18N-2.6P-10K (18-6-12) Osmocote/m³ (11 lb/yd³), 2.0 kg concentrated superphosphate/m³ (3.5 lb/yd³) and 0.9 kg each of dolomite, gypsum, limestone flour, and Micro-Max trace elements/m³ (1.5 lb/yd³). Commercial medium B consisted of peat moss, pumice, conifer bark and sand (4:4:1:1, v/v) amended with 1.77 kg each of dolomite and limestone flour/m³ (3.0 lb/yd³), 0.77 kg ureaformaldehyde/m³ (1.3 lb/yd³), 0.6 kg conc. superphosphate/m³ (1.0 lb/yd³), 0.4 kg FeSO₄/m³ (0.7 lb/yd³), 0.3 kg each of KNO₃ and Ca(NO₃)₂/m³ (0.5 lb/yd³), and 75 g FTE 503/m³ (2 oz/yd³).

Before planting, media samples were dried and air-filled pore space (AS) (2) and total porosity (TP) (9) were measured on duplicate samples. Electrical conductivity (EC) was measured with an RD-26 Solubridge and pH by glass electrode in a 1:3 medium-water (w/w) slurry.

Rooted cuttings of *Photinia x fraseri* Dress. and *Thuja occidentalis* L. 'Pyramidalis,' pyramidal arbovitae, were planted into 15.0 x 17.5 cm (#1) containers of potting mixes on March 24, 1983 and placed in an unheated, clear polyethylene-covered greenhouse with 5 plants/plot and 5 replications of each treatment (media) in randomized block design. Greenhouse covers were removed on June 1.

¹Received for publication February 19, 1985; in revised form August 20, 1985. Oregon State University Agric. Exp. Stn. Tech. Paper No. 7407. This research supported in part by a grant from the Taulman-Weiss Co., Atlanta, GA

²Professor, Associate Professor, and Research Technician, resp.

Table 1. Physical and chemical characteristics of potting media before planting.

Mix	Bulk density		EC	pH	AS	TP
	wet	dry				
	-----g/cm ³ -----					
			dS/m		% of total volume	
A	0.71	0.28	2.85	4.4	25	43
B	0.92	0.37	1.45	7.2	16	54
C ₁ P ₂ ^z	0.88	0.27	0.36	4.5	13	61
C ₂ P ₁	0.83	0.30	0.93	4.9	19	54
C ₁ B ₂	0.86	0.36	0.52	4.9	13	51
C ₂ B ₁	0.85	0.37	0.95	5.1	16	48
C ₁ S ₂	0.82	0.31	0.56	5.6	22	50
C ₂ S ₁	0.80	0.32	1.15	5.8	23	48
LSD (0.05)	0.04	0.03	0.22	0.2	5	4
Rate of compost ^y						
25%	0.85	0.31	0.48	5.0	16	54
50%	0.83	0.33	1.01	5.2	19	50
	* ^x	*	**	*	*	*
Organic component ^y						
Peat	0.85	0.29	0.65	4.6	16	58
Bark	0.85	0.37	0.74	5.0	15	50
Sawdust	0.81	0.32	0.86	5.7	23	49
LSD (0.05)	0.03	0.04	0.06	0.3	3	4

^zC = compost, P = peat, B = bark, S = sawdust, C₁P₂ = 25% (v/v) compost, 50% peat, 25% pumice.

^yMain effects for compost-based mixes only.

^x*, **: means significantly different at 1% and 5% levels, respectively.

All plants were irrigated daily by overhead sprinkler. Fertilizer was applied weekly as Peters 30N-4.3P-8.3K (30-10-10) (400 ppm N) from April 19 until June 15, when Osmocote 18N-2.6P-10K (18-6-12) was applied to all plants at 5 cc (1 tsp)/container. Leaf samples were taken at plant harvest on October 18, 1983 (*Thuja*) or October 28 (*Photinia*), washed in distilled water, and analyzed for elemental concentrations. Total N was determined by the Kjeldahl procedure. All other elements were determined by inductively-coupled argon plasma emission spectrometry (Plant Analysis Laboratory, Oregon State University) after dry ashing at 500 °C for 6 hr and extraction with 20% HNO₃. A composite sample of each medium was collected after plant harvest, dried, ground to pass a 20 mesh screen and analyzed for elemental concentrations as outlined for leaf analysis.

Results and Discussion

Increasing the proportion of compost in media from 25 to 50% increased initial dry bulk density, pH, EC, and AS but decreased TP and wet bulk density (Table 1). Compost-based media had higher bulk densities, pH and TP, and lower EC and AS than commercial medium A, but lower bulk density, pH and EC than commercial medium B. The organic component also affected media physical characteristics. Bulk densities were highest with bark; pH, EC and AS highest with sawdust; TP was highest with peat. These effects of

organic component on pH, EC, AS and TP of media are consistent with the physical and chemical properties of the individual organic components (5).

After 8 months of supporting plant growth, media properties varied significantly with the type of organic amendment. Shrinkage was significantly greater with peat as the organic amendment (Table 2). Dry bulk density and EC decreased, and pH increased with time, due in part to the high Ca and Mg content of the irrigation water (1.2 meq/L or 20 ppm). The physical and chemical properties of media did not vary significantly with proportion of compost.

In media samples taken at time of plant harvest, only Cu, Ni and Zn concentration appeared to increase consistently with increasing proportion of compost (Table 2). Levels of Al, Cd, Na, Ni, and Zn appeared to be higher in compost based media than in the commercial media.

Weight, height, and number of growing points were greatest for *Photinia* grown in medium B, followed by media containing peat and compost (Table 3). *Photinia* growth in medium A was limited because of early salt phytotoxicity from preincorporated fertilizer (Table 1).

Leaf N and P levels for *Photinia* grown in compost-based media were lower than in plants grown in medium A, but similar to levels for plants grown in medium B (Table 3). Leaf B concentration was consistently lower and leaf Zn concentration higher in compost-grown plants than in plants grown in the commercial media. Plant weight was negatively correlated with leaf N con-

tent ($r_{xy} = -0.70$, $P = 0.04$) in large part because of poor growth in medium A. Except for Zn, leaf trace element content did not increase appreciably with 50% compared to 25% compost. With the exception of B, leaf elemental concentrations were higher for *Photinia* in this experiment than were found in a recent survey of container-grown *Photinia* in commercial nurseries (R.L. Ticknor, unpublished data).

Medium A and the medium containing 50% compost and 25% peat produced the heaviest *Thuja* plants (Table 4). *Thuja* fresh weight was also greater in media containing 50 rather than 25% compost. Plant height was significantly increased with 50% compost only when sawdust was part of the medium. Proportion of compost in media had little effect on leaf tissue elemental concentrations, with the higher rate resulting only in small increases in Ni and Zn, and decreases in S, Fe and Mn. Levels of Cd, Mn, Ni, and Zn were higher in the compost:peat media, presumably due to the lower initial pH. There was no correlation between height, weight, and tissue nutrient concentration of plants grown in either commercial or compost media.

In a previous study with annual bedding plants, plant growth correlated significantly with high initial AS of the media (5). In contrast, *Photinia* and *Thuja* growth did not significantly correlate with any of the initial or final physical and chemical characteristics of the media. In every case, the effects on media characteristics of increasing the proportion of compost from 25 to 50% were the opposite of those associated with peat as organic component (Table 1). However, both the higher

rate of compost and presence of peat in the media favored plant growth. Better plant growth may have been more closely correlated with initial nutrient content or nutrient binding capacity of the compost (not measured). Chemical and physical characteristics of media measured before use may have little value in predicting woody plant growth, at least when AS, TP, pH, EC, etc. do not differ greatly from established ranges for acceptable growth (3,7,13). Maintaining a critical minimum AS and maximum TP was found essential for adequate growth of container-grown *Rhododendron*, in a case where addition of peat to the mix depressed growth due to excessively high water holding capacity (1).

The compost-based media exhibited considerable stability over time in terms of bulk density, AS, WS, and shrinkage (Tables 1 and 2). Composts which decompose rapidly or have extremely high EC or high pH resistant to change have reduced growth of several container-grown woody ornamentals (11).

Supplemental N was needed with all media. Leaf P levels of plants grown in compost-based media were generally comparable to those grown in the phosphate-supplemented commercial media (Tables 3 and 4). Based on these results, fertilizers with only low levels of phosphate would be needed to maintain adequate phosphate in the media. Potassium levels in leaves were adequate when liquid fed with 30N-4.3P-8.3K (30-10-10) and top dressed with Osmocote 18N-2.6P-10K (18-6-12). Potassium levels in the compost-based media analyzed at the end of the growing season were lower

Table 2. Physical characteristics and elemental concentration of media after eight months of plant growth.

Mix	Bulk Density		EC	pH	AS	TP	Shrinkage	N	P	K	Ca	Mg	S	Al	B	Cd	Cu	Fe	Mn	Mo	Na	Ni	Zn
	wet	dry																					
	- g/cm ³ -	- dS/m																					
A	0.77	0.21	1.40	6.8	26	56	9	0.58	0.11	0.21	1.40	0.24	0.17	1549	5	1.6	46	3256	137	0.7	117	3	53
B	0.91	0.30	0.18	7.6	15	46	13	0.81	0.02	0.06	0.53	0.09	0.07	1025	3	0.6	15	1627	64	0.4	166	1	14
C ₁ P ₂ ²	0.85	0.22	0.18	7.6	21	63	18	0.37	0.07	0.06	0.48	0.09	0.10	1629	3	1.5	29	2472	40	0.7	224	7	53
C ₂ P ₁	0.80	0.26	0.25	7.4	26	54	13	0.30	0.20	0.07	0.49	0.09	0.11	2535	3	4.5	76	4129	96	1.1	282	17	186
C ₁ B ₂	0.73	0.30	0.30	7.1	23	50	9	0.50	0.09	0.06	0.48	0.09	0.07	1694	3	2.0	35	2492	82	0.7	208	8	81
C ₂ B ₁	0.79	0.27	0.25	6.9	29	50	10	0.55	0.25	0.08	0.48	0.09	0.11	3051	4	5.7	73	3480	98	1.5	320	13	159
C ₁ S ₂	0.81	0.26	0.38	7.3	30	51	11	0.52	0.22	0.07	0.40	0.09	0.13	2449	3	3.6	48	2864	118	1.0	357	8	137
C ₂ S ₁	0.81	0.26	0.33	7.4	23	64	11	0.38	0.19	0.06	0.32	0.06	0.10	1985	3	3.6	53	2655	88	0.9	286	11	147
LSD (0.05)	0.05	0.05	0.30	0.4	6	7	2																
Rate of compost³																							
25%	0.80	0.26	0.29	7.3	25	55	13	0.46	0.13	0.06	0.45	0.09	0.10	1924	3	2.4	37	2609	80	0.8	263	8	90
50%	0.83	0.26	0.28	7.3	26	56	11	0.41	0.21	0.07	0.43	0.08	0.11	2524	3	4.6	67	3421	94	1.2	296	14	164
	NS ⁴	NS	NS	NS	NS	NS	NS																
Organic component³																							
Peat	0.83	0.24	0.21	7.5	23	59	15	0.34	0.13	0.07	0.49	0.09	0.10	2082	3	3.0	53	3300	68	0.9	253	12	120
Bark	0.76	0.29	0.27	7.0	26	50	10	0.52	0.17	0.07	0.48	0.09	0.09	2372	3	3.8	54	2986	90	1.1	164	11	120
Sawdust	0.81	0.26	0.35	7.4	26	58	11	0.45	0.21	0.07	0.36	0.08	0.12	2217	3	3.6	51	2760	103	1.0	322	10	142
LSD (0.05)	0.03	0.04	0.10	0.2	NS	6	3																

²C = compost, P = peat, B = bark, S = sawdust; C₁P₂ = 25% (v/v) compost, 50%, 25% pumice.

³Main effects for compost-based mixes only.

⁴NS: means not significantly different at 5% level.

Table 3. Growth and leaf elemental concentration of *Photinia* grown in compost-based and commercial media.

Mix	Weight	Height	No. of growing points	Leaf elemental concentrations												
				N	P	K	Ca	Mg	S	B	Cd	Cu	Fe	Mn	Ni	Zn
	(g)	(cm)		----- % -----						----- mg/kg -----						
A	88	44	20	2.13	0.40	1.6	1.1	0.23	0.20	21	0.14	5.8	258	43	1.0	45
B	157	51	45	1.82	0.23	1.4	1.2	0.28	0.15	29	0.08	3.2	276	32	0.7	26
C ₁ P ₂ ^z	129	50	23	1.72	0.23	1.4	1.0	0.23	0.17	13	0.12	5.2	248	34	1.6	46
C ₂ P ₁	150	55	29	1.59	0.29	1.4	1.1	0.24	0.17	15	0.15	5.6	226	38	2.4	74
C ₁ B ₂	90	41	24	1.91	0.32	1.6	1.1	0.24	0.16	17	0.13	4.4	245	33	1.5	60
C ₂ B ₁	122	50	30	1.73	0.31	1.5	1.1	0.23	0.16	17	0.18	5.8	274	35	1.9	87
C ₁ S ₂	86	43	23	1.92	0.27	1.6	1.0	0.23	0.21	15	0.19	4.4	210	39	1.2	70
C ₂ S ₁	101	47	23	1.89	0.30	1.7	1.2	0.24	0.19	17	0.19	4.2	240	41	1.2	90
LSD (0.05)	39	8	11	0.10	0.05	0.2	0.2	0.03	0.04	7	0.07	1.1	NS	NS	0.4	12
Rate of compost^y																
25%	102	45	23	1.85	0.27	1.5	1.0	0.23	0.18	15	0.15	4.7	234	35	1.4	59
50%	124	51	27	1.74	0.30	1.5	1.1	0.24	0.17	16	0.17	5.2	247	38	1.8	84
	** ^x	**	*	*	NS	NS	NS	NS	NS	NS	*	NS	NS	NS	NS	**
Organic component^y																
Peat	140	53	26	1.66	0.26	1.4	1.1	0.24	0.17	14	0.14	5.4	237	36	2.0	60
Bark	106	46	27	1.82	0.32	1.6	1.1	0.24	0.16	17	0.16	5.1	260	34	1.7	68
Sawdust	94	45	23	1.91	0.29	1.7	1.1	0.24	0.20	16	0.19	4.3	225	40	1.2	80
LSD (0.05)	10	4	3	0.11	0.04	0.2	NS	NS	0.03	NS	0.04	0.8	NS	NS	0.7	6

^zC = compost, P = peat, B = bark, S = sawdust, C₁P₂ = 25% (v/v) compost, 50% peat, 25% pumice.

^yMain effects for compost-based mixes only.

^x**, *, NS: mean significantly different at 1% and 5% levels, and nonsignificant, respectively.

than in the commercial media, indicating possible need for a K-containing fertilizer after potting. It may be necessary to add Ca and Mg at locations where the irrigation water contains low levels of these elements.

Addition of minor elements to compost-based media does not appear necessary, with some possible exceptions. Compost-based media tended to have low B concentration and leaves of *Photinia* grown in these media had lower B concentration than did plants in the commercial media. A spray application of B during the growing season might be necessary to prevent a deficiency with low-B composts. However, some composts contain high B levels (8).

Significance to the Nursery Industry

Sewage sludge composted with conifer sawdust in a closed vessel system is a suitable weed free component of media for growing container nursery stock. Plant growth in media containing 50% compost was equal to or better than that in two commercial media. One pound of slow release nitrogen per cubic yard would be the only fertilizer required initially with a 50% compost medium. If compost is priced competitively with other possible components, total costs for media and fertilizers would be reduced. Successful production requires maintaining appropriate nutrient, aeration, and moisture levels in the chosen medium. Irrigation and fertilizer applications will need to be altered when making changes in media components.

Literature Cited

1. Bilderback, T.C., W.C. Fonteno, and D.R. Johnson. 1982. Physical properties of media composed of peanut hulls, pine bark, and peat moss and their effects on azalea growth. J. Amer. Soc. Hort. Sci. 107:522-525.
2. Buscher, F.K. and D. Van Doren. 1973. Determination of air-filled pore space for container grown nursery stock. Proc. Intern. Plant Prop. Soc. 23:232-233.
3. DeBodt, M. and O. Verdonck. 1972. The physical properties of the substrates in horticulture. Acta Hort. 26:37-44.
4. Hemphill, D.D., Jr., T.L. Jackson, L.W. Martin, G.L. Kiemnec, D. Hanson, and V.V. Volk. 1982. Sweet corn response to application of three sewage sludges. J. Environ. Qual. 11:191-196.
5. Hemphill, D.D., Jr., R.L. Ticknor, and D.J. Flower. 1984. Growth response of annual transplants and physical and chemical properties of growing media as influenced by composted sewage sludge amended with organic and inorganic materials. J. Environ. Hort. 2:112-116.
6. Gouin, F.R. 1977. Conifer tree seedling response to nursery soil amended with composted sewage sludge. HortScience 12:341-342.
7. Joiner, J.N. and C.A. Conover. 1965. Characteristics affecting desirability of various media components for production of container-grown plants. Proc. Soil Crop Soc. Fla. 25:320-327.
8. Lumis, G.P. and A.G. Johnson. 1982. Boron toxicity and growth suppression of *Forsythia* and *Thuja* grown in mixes amended with municipal waste compost. HortScience 17:821-822.
9. Mazur, A.R., T.D. Hughes, and J.B. Gartner. 1975. Physical properties of hardwood bark growth media. HortScience 10:30-33.
10. McCoy, M. and J.L. Green. 1984. Composted sludge and granulated straw in container media produce good growth if used properly. Amer. Nurs. 159(1):73-75, 78-81.

11. Sanderson, K.C. 1980. Use of sewage refuse compost in the production of ornamental plants. HortScience 15:173-178.

12. Spomer, L.A. 1974. Optimizing container soil amendment: the "threshold proportion" and prediction of porosity. HortScience 9:532-533.

13. Waters, W.E., W. Llewellyn, and J. Nesmith. 1970. The chemical, physical, and salinity characteristics of twenty-seven soil media. Proc. Fla. State Hort. Soc. 83:482-488.

14. Wootton, R.D., F.R. Gouin, and F.C. Stark. 1981. Composted, digested sludge as a medium for growing flowering annuals. J. Amer. Soc. Hort Sci. 106:46-49.

Table 4. Growth and leaf elemental concentration of *Thuja* grown in compost-based and commercial media.

Mix	Weight	Height	Leaf elemental concentrations												
			N	P	K	Ca	Mg	S	B	Cd	Cu	Fe	Mn	Ni	Zn
	g	cm	-----%						-----mg/kg-----						
A	98	69	2.25	0.39	0.95	1.8	0.39	0.16	20	0.11	6.0	250	213	1.0	55
B	81	62	2.18	0.36	0.89	1.8	0.34	0.18	27	0.09	3.2	202	203	0.5	33
C ₁ P ₂ ^z	82	67	2.31	0.35	0.89	1.8	0.34	0.19	19	0.22	5.6	266	179	2.2	89
C ₂ P ₁	95	65	2.06	0.33	0.86	1.7	0.35	0.17	18	0.20	5.7	254	122	3.2	119
C ₁ B ₂	69	64	2.22	0.35	0.91	1.7	0.35	0.18	22	0.13	5.9	320	113	2.0	83
C ₂ B ₁	76	65	2.36	0.37	0.91	1.7	0.33	0.17	22	0.12	6.3	243	109	2.5	97
C ₁ S ₂	55	57	2.37	0.35	0.87	1.7	0.31	0.18	21	0.16	5.7	309	139	1.7	75
C ₂ S ₁	75	63	2.24	0.36	0.88	1.8	0.32	0.16	21	0.16	6.3	273	130	2.3	89
LSD (0.05)	13	5	0.16	0.03	0.04	0.1	0.02	NS	2	0.06	0.9	NS	38	0.5	15
Rate of compost^y															
25%	69	63	2.30	0.35	0.89	1.7	0.33	0.18	21	0.17	5.7	299	144	2.0	82
50%	82	64	2.22	0.35	0.89	1.7	0.33	0.17	20	0.16	6.0	257	120	2.6	102
	** ^x	NS	NS	NS	NS	NS	NS	*	NS	NS	NS	*	*	*	*
Organic component^y															
Peat	89	66	2.18	0.34	0.89	1.7	0.35	0.18	19	0.21	5.7	260	150	2.6	104
Bark	73	65	2.29	0.36	0.91	1.7	0.34	0.18	22	0.13	6.1	281	112	2.2	90
Sawdust	65	60	2.31	0.36	0.88	1.7	0.32	0.17	21	0.16	6.0	292	135	2.0	82
LSD (0.05)	10	NS	NS	NS	NS	NS	0.02	NS	NS	0.06	NS	NS	26	0.4	6

^zC = compost, P = peat, B = bark, S = sawdust; C₁P₂ = 25% (v/v) compost, 50% peat, 25% pumice.

^yMain effects for compost-based mixes only.

^x**, *, NS: means significantly different at 1% and 5% levels, and non-significant, respectively.